

The cross-correlation distribution of a p -ary m -sequence of period $p^{2m} - 1$ and its decimation by $\frac{(p^m+1)^2}{2(p^e+1)}$

Yuhua Sun¹, Hui Li¹, Zilong Wang¹

¹ State Key Laboratory of Integrated Service Networks,

Xidian University, Xi'an, 710071, Shaanxi, China

Email: sunyuhua-1@163.com lihui@mail.xidian.edu.cn

wzlmath@gmail.com

November 16, 2011

Abstract

Let $n = 2m$, m odd, $e|m$, and p odd prime with $p \equiv 1 \pmod{4}$. Let $d = \frac{(p^m+1)^2}{2(p^e+1)}$. In this paper, we study the cross-correlation between a p -ary m -sequence $\{s_t\}$ of period $p^{2m} - 1$ and its decimation $\{s_{dt}\}$. Our result shows that the cross-correlation function is six-valued and that it takes the values in $\{-1, \pm p^m - 1, \frac{1 \pm p^{\frac{e}{2}}}{2} p^m - 1, \frac{(1-p^e)}{2} p^m - 1\}$. Also, the distribution of the cross-correlation is completely determined.

Index Terms. finite field; p -ary m -sequence; decimation; cross-correlation;

1 Introduction

One problem of considerable interest has been to find a decimation value d such that the cross-correlation between a p -ary m -sequence $\{s_t\}$ of period $p^n - 1$ and its decimation $\{s_{dt}\}$ is low. When $\gcd(d, p^n - 1) = 1$, the decimated sequence $\{s_{dt}\}$ is also an m -sequence of period $p^n - 1$. Basic results on the cross-correlation between two m -sequences can be found in [1-3]. When $\gcd(d, p^n - 1) \neq 1$, the sequence $\{s_{dt}\}$ has period $\frac{p^n - 1}{\gcd(d, p^n - 1)}$. For this case, there also are many good results which can be found in [4-7].

In [6], for an odd prime p , Choi, Lim, No, and Chung investigated into the cross-correlation of a p -ary m -sequence of period $p^n - 1$ and its decimated sequence by $d = \frac{(p^m+1)^2}{2(p+1)}$, where $n = 2m$ and m is odd. They have shown that the magnitude of the cross-correlation values is upper bounded by $\frac{p+1}{2} p^{\frac{n}{2}} + 1$. Recently, for $n = 2m$, $p \equiv 3 \pmod{4}$, Luo, Helleseeth, and Kholosha [7] determined the distribution of the cross-correlation values of a p -ary m -sequence $\{s_t\}$ of period $p^n - 1$ and its decimated sequence $\{s_{dt}\}$ by $d = \frac{(p^m+1)^2}{2(p^e+1)}$, where m is odd and $e|m$. They derived that the cross-correlation is six-valued.

It is not difficult to observe that the decimation value d in [7] is a generalization of that in [6], i.e., for $p \equiv 3 \pmod{4}$ the result in [6] can be generalized to a more general form. In fact, it can also be generalized to the case of $p \equiv 1 \pmod{4}$. In this paper, for $p \equiv 1 \pmod{4}$, we will combine the machinery in [5],[7] and [6] to study the cross-correlation between a p -ary m -sequence $\{s_t\}$ of period $p^{2m} - 1$ and its decimated sequence $\{s_{dt}\}$ by the same d as that in [7], that is, $d = \frac{(p^m+1)^2}{2(p^e+1)}$, $e|m$ and m is odd. We show that the cross-correlation is also six-valued. And the distribution of the cross correlation values is determined.

2 Preliminaries

For an odd prime p , let F_{p^n} denote the finite field with p^n elements and $F_{p^n}^* = F_{p^n} \setminus \{0\}$. The trace function Tr_m^n from the field F_{p^n} onto the subfield F_{p^m} is defined by

$$\text{Tr}_m^n(x) = x + x^{p^m} + x^{p^{2m}} + \cdots + x^{p^{(h-1)m}},$$

where $h = n/m$.

We will use the following notations in the remaining part of this paper unless otherwise specified. Let $p \equiv 1 \pmod{4}$, $n = 2m$, $e|m$ and $d = \frac{(p^m+1)^2}{2(p^e+1)}$, where m is an odd integer. Let α be a primitive element of F_{p^n} . Then we have $\gcd(d, p^n - 1) = \frac{p^m+1}{2}$ and $d(p^{m+e} + 1) \equiv p^m + 1 \pmod{p^n - 1}$ (These also can refer to [6] or [7]). It should be pointed out that both $\frac{p^m+1}{2}$ and d are odd when $p \equiv 1 \pmod{4}$.

A p -ary m -sequence $\{s_t\}$ is given by

$$s_t = \text{Tr}_1^n(\alpha^t),$$

where Tr_1^n is the trace function from F_{p^n} onto F_p .

The periodic cross correlation function $C_d(\tau)$ between $\{s_t\}$ and $\{s_{dt}\}$ is defined by

$$C_d(\tau) = \sum_{t=0}^{p^n-2} \omega^{s_{t+\tau} - s_{dt}},$$

where $0 \leq \tau \leq p^n - 2$ and ω is a primitive complex p th root of unity.

Definition 1 ([8]) *A quadratic form $x_1, x_2, \dots, x_n$ in n indeterminates over F_p is a homogeneous polynomial in $F_p[x_1, x_2, \dots, x_n]$ of degree 2, that is,*

$$f(x_1, x_2, \dots, x_n) = \sum_{i,j=1}^n a_{i,j} x_i x_j \quad \text{with } a_{i,j} = a_{j,i} \in F_p.$$

The $n \times n$ matrix A whose (i, j) entry is $a_{i,j}$ is called the coefficient matrix of f . From [8], we know that every quadratic form over F_p is equivalent to a diagonal quadratic form $a_1x_1^2 + a_2x_2^2 + \cdots + a_rx_r^2$ over F_p , where $r \leq n$ is called the rank of f . For any $b \in F_p$, the number of solutions of $f(x_1, x_2, \dots, x_n) = b$ is equal to the number of solutions of $a_1x_1^2 + a_2x_2^2 + \cdots + a_rx_r^2 = b$. We denote $\Delta = a_1a_2 \cdots a_r$, then Δ is called the determinant of f .

Definition 2 ([8]) *The quadratic character of F_{p^n} is defined as*

$$\eta(x) = \begin{cases} 1, & \text{if } x \text{ is a nonzero square in } \text{GF}(p^n) \\ -1, & \text{if } x \text{ is a nonsquare in } \text{GF}(p^n) \\ 0, & \text{if } x = 0. \end{cases}$$

Definition 3 ([8]) *The canonical additive of F_{p^n} is defined as*

$$\chi(x) = \sum_{x \in F_{p^n}} \omega^{\text{Tr}_1^n(x)}.$$

Definition 4 ([8]) *The Gaussian sum $G(\eta, \chi)$ of η and χ is defined as*

$$G(\eta, \chi) = \sum_{x \in F_{p^n}^*} \eta(x)\chi(x).$$

3 The ranks of two quadratic forms

In this section, we will give some results to prove our main theorem. First, using the similar method of [6] or [7], we can get the following four lemmas.

Lemma 1 *Let the symbols be defined as in section 2. Then the cross-correlation between $\{s_t\}$ and $\{s_{dt}\}$ is given by*

$$\begin{aligned} C_d(\tau) &= -1 + C(-1, c) = -1 + \sum_{x \in F_{p^n}} \chi(-x^d + cx) \\ &= -1 + \frac{1}{2} (E(-1, c) + E(-\alpha^d, c\alpha)) \end{aligned}$$

where $c = \alpha^\tau$, and $E(a, b) = \sum_{x \in F_{p^n}} \chi(ax^{p^m+1} + bx^{p^{m+e}+1})$. Further, $q_{a,b}(x) = \text{Tr}_1^n(ax^{p^m+1} + bx^{p^{m+e}+1})$ is a quadratic form over F_p . □

Let $r_{a,b}$ be the rank of the quadratic form $q_{a,b}(x)$. Then the following Lemma can be derived from Corollary 5 of [6].

Lemma 2 Let $p \equiv 1 \pmod{4}$. Let Δ is the determinant of $q_{a,b}(x)$, where $x \in \mathbb{F}_{p^n}$. Then we have

$$E(a, b) = \sum_{x \in \mathbb{F}_p^n} \omega^{q_{a,b}(x)} = \pm p^{n - \frac{r_{a,b}}{2}},$$

where $\pm$ depends on $\eta(\Delta)$ and η is the quadratic character of $\mathbb{F}_p$. □

By Lemmas 1 and 2, in order to compute $C_d(\tau)$, we need to find the values of $r_{-1,c}$ and $r_{-\alpha^d, c\alpha}$. The following Lemma which can be found in [6] or [4] gives us a method to compute them.

Lemma 3 ([6],[4]) Let $f(x) \in \mathbb{F}_{p^n}[x]$ can be expressed as a quadratic form in $\text{GF}(p)[x_1, x_2, \dots, x_n]$, where $x \in \mathbb{F}_{p^n}$. Furthermore, let

$$Y = \{y \in \mathbb{F}_{p^n} : f(x+y) = f(x) \text{ for all } x \in \mathbb{F}_{p^n}\}.$$

Then $\text{rank}(f) = n - \text{Log}_p |Y|$. □

Using Lemma 3 and the method of [6] or [7], we can get the following lemma.

Lemma 4 Let the symbols be defined as before. Then the number of solutions of $q_{a,b}(x+y) = q_{a,b}(x)$ for all $x \in \mathbb{F}_{p^n}$ is equal to the number of solutions of

$$b^{p^{m+e}} y^{p^{2e}} + (a^{p^{m+e}} + a^{p^e}) y^{p^e} + by = 0 \quad (1)$$

in $\mathbb{F}_{p^n}$. Furthermore, Since Eq.(1) is a $\mathbb{F}_{p^e}$ -linearized polynomial, we know that the number of the solutions of it is 1, p^e or p^{2e} . Hence, $r_{a,b} = n, n-e$ or $n-2e$. □

In fact, we can get further result about $E(-\alpha^d, c\alpha)$. To this end, we need the following lemmas and corollary.

Lemma 5 (Theorem 5.30 of [8]) Let the symbols be defined as in section 2. Then

$$\sum_{x \in \mathbb{F}_{p^n}} \chi(ax^2) = \eta(a)G(\eta, \chi),$$

where $a \in \mathbb{F}^*$. □

Lemma 6 (Theorem 5.16 of [8]) Let the symbols be defined as in section 2. Then

$$G(\eta, \chi) = \begin{cases} p^m & \text{if } \frac{p^m+1}{2} \text{ even,} \\ -p^m & \text{if } \frac{p^m+1}{2} \text{ odd.} \end{cases}$$

□

Corollary 1 *Let the symbols be defined as in section 2. Then we have*

$$\sum_{x \in F_{p^n}} \chi(ax^2) = -\eta(a)p^m,$$

where $a \in F^*$.

Proof: Note that $\frac{p^m+1}{2}$ is odd when $p \equiv 1 \pmod{4}$. Combining Lemmas 5 and 6, the result follows. $\square$

Lemma 7 *Let the symbols be defined as before. Then*

$$(1) E(-\alpha^d, c\alpha) = \eta(c)p^m.$$

$$(2) r_{-\alpha^d, c\alpha} = n.$$

Proof: (1) Similar to the proof of Lemma 1 in [7], we can get

$$E(-\alpha^d, c\alpha) = \sum_{x \in F_{p^n}} \chi(cx^2).$$

Note that α is a nonsquare in F_{p^n} . By Corollary 1, we can get

$$E(-\alpha^d, c\alpha) = -\eta(c\alpha)p^m = (-1)\eta(\alpha)\eta(c)p^m = \eta(c)p^m.$$

(2) By lemma 2 and the above result, we can get that $r_{-\alpha^d, c\alpha} = n$. $\square$

Combining Lemmas 1, 2, 4 and 7, we can get the following corollary.

Corollary 2 *Let the symbols be defined as above. Then we have*

$$E(-1, c) \in \{\pm p^m, \pm p^{m+\frac{e}{2}}, \pm p^{m+e}\}, E(-\alpha^d, c\alpha) \in \{\pm p^m\}$$

and

$$C(-1, c) \in \{0, \pm p^m, \frac{1 \pm p^{\frac{e}{2}}}{2}p^m, \frac{-1 \pm p^{\frac{e}{2}}}{2}p^m, \frac{\pm(1-p^e)}{2}p^m, \frac{\pm(1+p^e)}{2}p^m\}.$$

$\square$

In fact, we can prove $C(-1, c) \neq \frac{-1 \pm p^{\frac{e}{2}}}{2}p^m$. Before proving it, we need the following lemma.

Lemma 8 ([9]) *Let $g_v(z) = z^{p^e+1} - vz + v$, where $v \in F_{p^n}^*$. Then the following results hold.*

(1) *The Equation $g_v(z) = 0$ has either 0, 1, 2, or $p^e + 1$ roots in F_{p^n} .*

(2) *If $g_v(z) = 0$ has only one root in F_{p^n} , then c is a square element in F_{p^n} ;*

(3) *Let N_1 denote the number of $v \in F_{p^n}^*$ such that $g_v(z) = 0$ has only one root in F_{p^n} . Then $N_1 = p^{n-e}$.*

(4) *If $g_v(z) = 0$ has only one root z_0 in F_{p^n} , then z_0 satisfies $(z_0 - 1)^{\frac{p^n-1}{p^e-1}} = 1$, i.e., $z_0 - 1$ is a $(p^e - 1)$ th power in F_{p^n} .* $\square$

Lemma 9 *Let the symbols be defined as before. Then the following two results hold.*

- (1) *If $r_{-1,c} = n - e$, then c is a square in $F_{p^n}^*$ and $E(-\alpha^d, c\alpha) = p^m$. Consequently, we get $C(-1, c) \neq \frac{-1 \pm p^{\frac{e}{2}}}{2} p^m$.*
- (2) *Let N_e be the number of c 's such that $r_{-1,c} = n - e$, where $c \in F_{p^n}^*$. Then $N_e = p^{m-e}(p^m + 1)$.*

Proof: (1) Let $f_c(x) = c^{p^{m+e}} x^{p^{2e}} - 2x^{p^e} + cx$, $y = x^{p^e-1}$ and $z = \frac{2}{c}y$. By Lemma 4, for $r_{-1,c} = n - e$, we know that the equation $f_c(x) = 0$ has $p^e - 1$ nonzero solutions in $F_{p^n}^*$ and that any two such solutions x_1, x_2 satisfy $x_1^{p^e-1} = x_2^{p^e-1}$, i.e., $g_c(y) = c^{p^{m+e}} y^{p^e+1} - 2y + c = 0$ has only one solution y_0 and y_0 is a $(p^e - 1)$ th power in $F_{p^n}^*$. Further, $g_c(y) = 0$ has only one solution in $F_{p^n}^*$ if and only if $g_v(z) = 0$ has only one solution $z_0 \in F_{p^n}^*$, where $v = \frac{4}{c(p^{m+1})p^e}$. Note that $c = \alpha^\tau$ and that $\beta = \alpha^{p^{m+1}}$ is a primitive element in F_{p^m} . Then we have $c^{p^{m+1}} = \beta^\tau$ and $v = 4\beta^{-\tau p^e} \in F_{p^m}^*$. By the same argument as that of Lemma 4 in [7], we can get that $g_v(z) = 0$ has only one root in $F_{p^n}^*$ if and only if $g_v(z) = 0$ has only one root in $F_{p^m}^*$. By Lemma 8 (2), when $g_v(z) = 0$ has only one root in $F_{p^m}^*$ the element v is a square in $F_{p^m}^*$, which implies that τ is even and that c is a square in F_{p^n} .

(2) Since $v = \frac{4}{c(p^{m+1})p^e} = \left(\frac{2}{c^{p^e}}\right)^{p^{m+1}} \in F_{p^m}^*$, then v runs through $p^m + 1$ times all the elements in $F_{p^m}^*$ when c runs through all the nonzero element in $F_{p^{2m}}^*$. By Lemma 8 (3), there are p^{m-e} v 's such that $g_v(z) = 0$ has only one root in $F_{p^m}^*$ when v runs through all the elements in $F_{p^m}^*$. Hence, there are $p^{m-e}(p^m + 1)$ c 's in $F_{p^{2m}}^*$ such that $g_v(z) = 0$ has only one root in $F_{p^m}^*$. By the argument of (1), there also are $p^{m-e}(p^m + 1)$ c 's such that $g_c(y) = 0$ has only one root in $F_{p^n}^*$. Let c_0 be any one of $p^{m-e}(p^m + 1)$ c 's and let $y_0 \in F_{p^{2m}}^*$ be the only root of $g_{c_0}(y) = 0$. Next, we will prove that y_0 must be a $(p^e - 1)$ th power in $F_{p^{2m}}^*$, i.e., $y_0^{\frac{p^{2m}-1}{p^e-1}} = 1$. Let $z_0 = \frac{2}{c_0}y_0 \in F_{p^m}^*$. Then $z_0^{p^e+1} - \frac{4}{c_0(p^{m+1})p^e}z_0 + \frac{4}{c_0(p^{m+1})p^e} = 0$, i.e.,

$$z_0 - 1 = \frac{c_0^{(p^m+1)p^e} z_0^{p^e+1}}{4}. \text{ By Lemma 8 (4), } (z_0 - 1)^{\frac{p^m-1}{p^e-1}} = \left(\frac{c_0^{(p^m+1)p^e} z_0^{p^e+1}}{4}\right)^{\frac{p^m-1}{p^e-1}} = \left(\frac{c_0^{p^m+1} z_0^2}{4}\right)^{\frac{p^m-1}{p^e-1}} = \left(\frac{c_0^{p^m+1} (\frac{2}{c_0} y_0)^2}{4}\right)^{\frac{p^m-1}{p^e-1}} = \left(c_0^{p^m-1} y_0^2\right)^{\frac{p^m-1}{p^e-1}} = 1, \text{ which implies that there exists an element } \theta \in F_{p^m} \text{ such}$$

$$\text{that } c_0^{p^m-1} y_0^2 = \theta^{p^e-1}, \text{ i.e., } y_0^{\frac{p^n-1}{p^e-1}} = \left[\frac{\theta^{p^e-1}}{c_0^{p^m-1}}\right]^{\frac{p^n-1}{2(p^e-1)}} = \frac{\theta^{(p^m-1) \cdot \frac{p^m+1}{2}}}{c_0^{\frac{p^m-1}{p^e-1} \cdot \frac{p^n-1}{2}}} = \frac{1}{\left[\frac{c_0^{\frac{p^n-1}{2}}}{c_0^{\frac{p^n-1}{2}}}\right]^{\frac{p^m-1}{p^e-1}}} = 1. \text{ By the result}$$

of (1), we know that c_0 is a square in F_{p^n} , then $c_0^{\frac{p^n-1}{2}} = 1$. Hence, we have $y_0^{\frac{p^n-1}{p^e-1}} = 1$, i.e., y_0 is a $(p^e - 1)$ th power in F_{p^n} . The result follows. $\square$

In the next section, we will find the equations which are satisfied by $E(-1, c)$ and $C(-1, c)$ in order to determine the distribution of $C(-1, c)$ or $C_d(\tau)$.

4 The distribution of $C_d(\tau)$

First, using the method of [5], we can get the following Lemma.

Lemma 10 *Let the symbols be defined as above. We have the following results:*

- (1) $\sum_{c \in F_{p^{2m}}} E(-1, c) = p^{2m}; E(-1, 0) = -p^m;$
- (2) $\sum_{c \in F_{p^{2m}}} [E(-1, c)]^2 = (2p^{2m} - 1)p^{2m};$
- (3) $\sum_{c \in F_{p^{2m}}} C(-1, c) = p^{2m}; C(-1, 0) = \frac{1}{2}(p^m - 1)p^m;$
- (4) $\sum_{c \in F_{p^{2m}}} [C(-1, c)]^2 = p^{4m}.$

□

Corollary 3 *Let $N_i = |\{c \in F_{p^n}^* | r_{-1, c} = n - i\}|$, where $i = 0, e, 2e$. Then we have*

$$N_0 = \frac{(p^{m+2e} - p^{m+e} - p^m - p^{2e} + 2)(p^m + 1)}{p^{2e} - 1},$$

$$N_e = p^{m-e}(p^m + 1),$$

$$N_{2e} = \frac{(p^{m-e} - 1)(p^m + 1)}{p^{2e} - 1}.$$

Proof: By Lemma 9 (2), we know that $N_e = p^{m-e}(p^m + 1)$. Note that $N_0 + N_e + N_{2e} = p^{2m} - 1$. Further, by Lemma 10 (2), we have

$$N_0 + N_e p^e + N_{2e} p^{2e} = 2(p^{2m} - 1).$$

Straightforward calculation gives the result.

□

Combining the machinery in [5], [7] and [6], we have the following results.

Lemma 11 *Let the symbols be defined as above. Then*

- (1) $\sum_{c \in F_{p^{2m}}} [E(-1, c)]^3 = (-p^{2m} + p^{m+e} + p^e)p^{3m}.$
- (2) $\sum_{c \in F_{p^{2m}}} [C(-1, c)]^3 = \frac{1}{8}p^{3m}(p^{3m} - p^{2m} + p^{m+e} + 6p^m + p^e).$

Corollary 4 Let $N_{i,\epsilon} = |\{c \in \mathbb{F}_{p^n}^* | E(-1, c) = \epsilon p^{m+\frac{i}{2}}\}|$, where $i = 0, e, 2e$ and $\epsilon = \pm 1$. Then we have

$$\begin{aligned} N_{0,1} &= \frac{(p^{m+e} - 1)(p^m + 1)}{2(p^e + 1)}, \quad N_{0,-1} = \frac{(p^{m+e} - 2p^m - 2p^e + 3)(p^m + 1)}{2(p^e - 1)}, \\ N_{e,1} &= N_{e,-1} = \frac{1}{2}p^{m-e}(p^m + 1), \\ N_{2e,1} &= 0, \quad N_{2e,-1} = \frac{(p^{m-e} - 1)(p^m + 1)}{p^{2e} - 1}. \end{aligned}$$

Proof: Using Corollary 3, Lemma 10 (1) and Lemma 11 (1), we know that

$$\begin{aligned} N_{0,1} + N_{0,-1} &= N_0 = \frac{(p^{m+2e} - p^{m+e} - p^m - p^{2e} + 2)(p^m + 1)}{p^{2e} - 1}, \\ N_{e,1} + N_{e,-1} &= N_e = p^{m-e}(p^m + 1), \\ N_{2e,1} + N_{2e,-1} &= N_{2e} = \frac{(p^{m-e} - 1)(p^m + 1)}{p^{2e} - 1}. \end{aligned}$$

$$(N_{0,1} - N_{0,-1}) + p^{\frac{e}{2}}(N_{e,1} - N_{e,-1}) + p^e(N_{2e,1} - N_{2e,-1}) = p^m + 1, \quad (2)$$

$$(N_{0,1} - N_{0,-1}) + p^{\frac{3e}{2}}(N_{e,1} - N_{e,-1}) + p^{3e}(N_{2e,1} - N_{2e,-1}) = (p^m + 1)(-p^m + p^e + 1). \quad (3)$$

Note that both $p^{\frac{e}{2}}$ and $p^{\frac{3e}{2}}$ are irrational numbers. Eqs. (2) and (3) imply that $N_{e,1} = N_{e,-1}$. Straightforward calculation gives the result. $\square$

Theorem 1 Let $n = 2m$, $e|m$, where m is odd. Let $p \equiv 1 \pmod{4}$ and $d = \frac{(p^m - 1)^2}{2(p^e + 1)}$. Then we get the distribution of the cross correlation $C_d(\tau)$ in the following.

-1	occurs	$\frac{(p^{m+e} - 2p^m - 2p^e + 3)(p^m + 1)}{2(p^e - 1)}$	times
$p^m - 1$	occurs	$\frac{(p^e - 1)(p^m + 1)^2}{4(p^e + 1)}$	times
$-p^m - 1$	occurs	$\frac{p^{2m} - 1}{4}$	times
$\frac{1 \pm p^{\frac{e}{2}}}{2}p^m - 1$	occurs	$\frac{1}{2}p^{m-e}(p^m + 1)$	times
$\frac{1 - p^e}{2}p^m - 1$	occurs	$\frac{(p^{m-e} - 1)(p^m + 1)}{p^{2e} - 1}$	times

Proof: For convenience, we denote

$$N_{i,\epsilon_1,\epsilon_2} = |\{c \in \mathbb{F}_{p^{2m}}^* | E(-1, c) = \epsilon_1 p^{m+\frac{i}{2}} \text{ and } E(-\alpha^d, c\alpha) = \epsilon_2 p^m\}|,$$

where $i \in \{0, e, 2e\}$, $\epsilon_1 = \pm 1$, and $\epsilon_2 = \pm 1$. By Lemma 9 (1), we know that

$$N_{e,1,-1} = N_{e,-1,-1} = 0. \quad (4)$$

On the other hand, by Corollary 4, we know that

$$N_{e,1} = N_{e,1,1} + N_{e,1,-1} = N_{e,-1} = N_{e,-1,1} + N_{e,-1,-1} = \frac{1}{2}p^{m-e}(p^m + 1),$$

then we can get

$$N_{e,1,1} = N_{e,-1,1} = \frac{1}{2}p^{m-e}(p^m + 1). \quad (5)$$

Again, by Corollary 4, we can get

$$N_{0,1,1} + N_{0,1,-1} = N_{0,1} = \frac{(p^{m+e} - 1)(p^m + 1)}{2(p^e + 1)}, \quad (6)$$

$$N_{0,-1,1} + N_{0,-1,-1} = N_{0,-1} = \frac{(p^{m+e} - 2p^m - 2p^e + 3)(p^m + 1)}{2(p^e - 1)}, \quad (7)$$

$$N_{2e,1,1} + N_{2e,1,-1} = N_{2e,1} = 0, \quad (8)$$

$$N_{2e,-1,1} + N_{2e,-1,-1} = N_{2e,-1} = \frac{(p^{m-e} - 1)(p^m + 1)}{p^{2e} - 1} \quad (9)$$

Further, by Lemma 10 (3),(4) and Lemma 11 (2), we can get

$$\begin{aligned} (N_{0,1,1} - N_{0,-1,-1}) + \frac{1-p^e}{2}N_{2e,-1,1} - \frac{1+p^e}{2}N_{2e,-1,-1} \\ = \frac{1}{2}(-p^{m-e} + 1)(p^m + 1), \end{aligned} \quad (10)$$

$$\begin{aligned} (N_{0,1,1} + N_{0,-1,-1}) + \left(\frac{1-p^e}{2}\right)^2 N_{2e,-1,1} + \left(\frac{1+p^e}{2}\right)^2 N_{2e,-1,-1} \\ = \frac{1}{4}(2p^m - p^{m-e} - 1)(p^m + 1) \end{aligned} \quad (11)$$

$$\begin{aligned} (N_{0,1,1} - N_{0,-1,-1}) + \left(\frac{1-p^e}{2}\right)^3 N_{2e,-1,1} - \left(\frac{1+p^e}{2}\right)^3 N_{2e,-1,-1} \\ = \frac{1}{8}(2p^m + p^e + 1)(p^m + 1). \end{aligned} \quad (12)$$

Combining Eqs. (4.40-4.48), we can get the result. $\square$

Remark 1 For $p = 5$ and $e = 1$ in Theorem 1, the magnitude of the cross-correlation is upper bounded by $2\sqrt{p^n} + 1$. This is meaningful in CDMA communication systems.

References

- [1] T. Helleseeth, "Some results about the cross-correlation function between two maximal linear sequences," *Discrete Math.*, 16: 209-232, 1976
- [2] T. Helleseeth. Kumar P V. Sequences with low correlation, Chapter in: Handbook in Coding Theory. (Eds. V.S. Pless and W.C. Huffman), Elsevier 1998.
- [3] H. Dobbertin, T. Helleseeth, P.V. Kumar, and H. Martinsen, "Ternary m -sequences with three-valued cross-correlation function:new decimations of Welch and Niho type," *IEEE Trans. Inf. Theory*, vol. 47, no. 4, pp. 1473-1481, May 2001.
- [4] E. N. Müller, "On the cross-correlation of sequences over $GF(p)$ with short periods," *IEEE Trans. Inf. Theory*, vol.45, no. 1, pp. 289-295, Jan. 1999.
- [5] E.Y. Seo, Y.S. Kim,J.S. No, D. J. Shin, "Cross-Correlation Distribution of p -ary m -Sequence of Period $p^{4k} - 1$ and Its Decimated Sequences by $(\frac{p^{2k}+1}{2})^2$," *IEEE Trans. Inf. Theory*, vol.54, no. 7, pp. 3140-3149, July. 2008.
- [6] S.-T. Choi, J.-S. No, H. Chung,"On the cross-correlation of a p -nary m -sequence of period $p^{2m} - 1$ and its decimated sequence by $\frac{(p^m+1)^2}{2(p+1)}$," accepted by *IEEE Trans. Inf. Theory*.
- [7] J. Luo, T. Helleseeth, A. Kholosha,"Two Nonbinary Sequences with Six-valued Cross-Correlation" *Proceedings of IWSDA '11*, pp.44-47.
- [8] R. Lidl and H. Niederreiter, Finite fields, vol. 20 of Encyclopedia of Mathematics and Its Applications. Reading, MA: Addison-Wesley,1983.
- [9] Bluher, A.W.: On $x^{q+1} + ax + b$. Finite Fields Appl.10(3),285-305(2004)
- [10] T. Storer, Cyclotomy and Difference Sets, Lectures in Advanced Mathematics. Chicago, IL: Markham, 1967.